

CLASSIFICATION OF PLEISTOCENE DEPOSITS, NEW YORK CITY AND VICINITY--FULLER (1914) REVIVED AND REVISED

John E. Sanders and Charles Merguerian
Geology Department, Hofstra University
Hempstead, NY 11549-1140

INTRODUCTION

After the Pleistocene deposits in New York City and vicinity had been recognized as products of continental glaciation, debates arose and have continued over the number of times such now-vanished continental glacier(s) invaded our region. Late in the nineteenth- and early in the twentieth centuries, several authors cited evidence that SE New England had been subjected to at least two and probably three discrete ice advances (Shaler, 1866, 1886; Shaler, Woodworth, and Marbut, 1896; Upham, 1879a, b; 1880; 1889a, b, c; Woodworth, 1901; Clapp, 1908; Fuller, 1901, 1903, 1905, 1906). In the face of the case made by T. C. Chamberlin (1883, 1888, 1895a, b) for a "one-glacier-did-it-all" view, however, the multi-glaciation concept lost favor. The one-glacier view took over. (For examples, see references by Salisbury and co-workers in New Jersey.)

In 1914, the U. S. Geological Survey finally published M. L. Fuller's monograph on the geology of Long Island (based on field work completed nearly a decade earlier). Fuller found deposits that he interpreted as products of 4 glacial advances; between some of the glacial sediments, he found nonglacial strata. In order of decreasing age, Fuller (1914) recognized the following six major units of the Pleistocene on Long Island: (1) the Mannelto Gravel, (2) the Jameco Gravel, (3) the Gardiners Clay, (4) the Jacob Sand, (5) the Manhasset Formation (which Fuller diagnosed as consisting of two units of outwash, the basal Herod Gravel and upper Hempstead Sand/Gravel, that are separated by a middle Montauk Till) and (6) the two world-famous terminal-moraine ridges and associated outwash plains to the south of each: the older Ronkonkoma and younger Harbor Hill, which overlie the Manhasset Formation (Table 1).

In 1934, Woodworth and Wigglesworth published their monograph on the glacial deposits of Cape Cod and offshore islands. Their stratigraphic units closely matched those Fuller had found on Long Island; they accepted and reinforced Fuller's four-glacier interpretation. Almost immediately thereafter, revisionist-minded "Friends of the Pleistocene" (Flint, 1935; Fleming, 1935; MacClintock and Richards, 1936, 1937) cast aside the four-glacier classification of Fuller and of Woodworth and Wigglesworth and adopted the viewpoint that nearly all of the glacial features in our region had been made during the latest glacial episode, the "Wisconsinan" of modern terminology. This was, in effect, a return to T. C. Chamberlin's interpretation. The contemporary view among virtually all Pleistocene geologists (Schafer and Hartshorn, 1965; Muller, 1965; Koteff, Oldale, and Hartshorn, 1967; Sirkin, 1968; 1982, 1986, 1991; Sirkin and Mills, 1975; Cadwell, 1986, 1989) widely circulated in textbooks, discussed at conferences, and in popular accounts of the geology of Long Island (Schuberth, 1968; Isachsen, Landing, Lauber, Rickard, and Rogers, 1991) and of New Jersey (Widmer, 1964), is that the latest glacier to visit our area, i. e., the "Woodfordian" glacier, deposited not only both of Long Island's terminal-moraine ridges but also produced all the other glacial features in the area. In

short, they support the view that one glacier did it all. The one-glacier view had become so firmly entrenched that evidence to the contrary was cited as constituting the "two-till **problem**" (Pessl and Schafer, 1968).

But, despite the continued onward rolling of the "one-glacier" bandwagon, several investigators have published evidence comparable to that mentioned in a previous paragraph in support of a multi-glacial interpretation (Goldsmith, 1960; Flint, 1961; Kaye, 1964b, c, d; 1976, 1982; Pessl, 1966; Oldale, 1982; Oldale and Colman, 1992; Oldale and Eskenasy, 1983; Oldale, Valentine, Cronin, Spiker, Blackwelder, Belknap, Wehmiller, and Szabo, 1981; W. A. Newman and Mickelson, 1994; Muller and Calkin, 1993; Muller, Sirkin, and Craft, 1993).

Stage	Fuller, 1914	Fleming, 1935	MacClintock and Richards, 1936	Donner, 1964
Wisconsin	Harbor Hill Moraine Ronkonkoma Moraine	Harbor Hill Moraine Ronkonkoma Moraine	Harbor Hill Moraine Ronkonkoma Moraine	Harbor Hill Moraine Ronkonkoma Moraine
		Hempstead Gravel	Manhasset Formation	Manhasset Fm. { Hempstead Gravel Montauk Till Herod Gravel
		Montauk Till		
		Herod Gravel		
Sangamon	Vineyard erosion surface		Jacob Sand Gardiners Clay	Gardiners Fm. { Jacob Sand Gardiners Clay
Illinoian	Hempstead Gravel } Montauk Till } Herod Gravel } Manhasset Fm.			
Yarmouth	Jacob Sand Gardiners Clay	Jacob Sand Gardiners Clay	Gardiners Clay (in part)	
Kansan	Jameco Gravel			
Pre-Kansan	Manetto Gravel			

Table 1. Pleistocene formations of Long Island showing Fuller's classification and two mid-1930's modifications to it proposed by revisionist-minded "Friends of the Pleistocene." In the "Fuller, 1914" column of Muller's original version of this table, the Harbor Hill Moraine and Ronkonkoma Moraine were shown as latest Wisconsin, the revisionist view. Fuller, however, assigned these two moraines to the Early Wisconsin, as shown here. (After E. H. Muller, 1965, tab. 2, p. 104.)

One of our chief contributions to knowledge of the local Quaternary record is an emphasis on the features by which the direction of flow of a former glacier can be established. We follow the principle that a continental glacier can be characterized locally by a distinctive flow direction. Application of this principle has led us to resurrect, but to modify in several important respects, M. L. Fuller's (1914) four-glacier classification of the Quaternary deposits of Long Island.

We have found five main reasons for resurrecting- and modifying Fuller's (1914) classification: (1) provenance data indicate that the latest-Wisconsinan ("Woodfordian" of

authors) glacier, which flowed from NNE to SSW, **did not reach most of eastern Long Island** but deposited a terminal moraine along the S coast of CT (Ellis, 1962; Flint and Gebert, 1976) and thus was not responsible for depositing the main body of the feature known as the Harbor Hill Moraine on Long Island (Fig. 1); (2) provenance- and structural data support the interpretation that the glacier which did deposit the main body of the Harbor Hill Moraine on




Figure 1. Sketch map of southern New England and vicinity, including adjacent islands showing locations of end-moraine ridges (black). We think that only the Old Saybrook and related moraines are products of the "Woodfordian" glacier that flowed from NNE to SSW. According to us, this moraine (of Glacier I in our proposed classification) was limited to coastal Connecticut as far W as the Norwalk Islands and extended southward so as to cross western Long Island (Queens and Brooklyn) only SW of these islands. (R. F. Flint and J. A. Gebert, 1974, fig. 4, p. 187.)

Long Island flowed regionally from NNW to SSE (Fig. 2) and was of pre-"Woodfordian" age (we offer no new data but accept Fuller's "Early Wisconsinan" assignment); (3) subsurface data from the vicinity of Jones Beach (Rampino, 1978 ms.; Rampino and Sanders, 1981) suggest an "Illinoian" age for the glacier that deposited the Ronkonkoma Moraine (earlier than Fuller's "Early Wisconsin"), and Gilbert-type deltas (G. K. Gilbert, 1885, 1890) in the Herod Gravel Member of Fuller's Manhasset Formation (Sanders, Merguerian, and Mills, 1993; Sanders and Merguerian, 1994b) that not only absolutely destroys the basis for the revisionists' "Wisconsinan" age reclassification of Fuller's Manhasset Formation but also requires Fuller's non-lacustrine paleoenvironmental analysis to be significantly modified by including a feature required for impounding a large Proglacial Lake Long Island (we think this feature probably was a now-eroded, pre-Ronkonoma terminal moraine but admit that it could have been an upbulged part of what is now the sea floor that was elevated as a peripheral bulge but disappeared after the

ice had melted); (4) the new-, and to us convincing, chronostratigraphic information that supports Fuller's "Yarmouthian" age of the Gardiners Clay that H. C. Ricketts (1986) collected from two borings made near Kings Point (on Great Neck, west side of Manhasset Bay) on which J. H. Wehmiller, of the University of Delaware, found D/L leucene values between 0.26 and 0.34, which implies that the age of the shells is about 225,000 years [225 Ka]; and (5) two much-weathered tills, not yet dated but presumably of Early Pleistocene ages, that we have discovered: a younger one at Croton Point, Westchester Co., NY; and an older one at Garvies Point, Long Island; and in SW Staten Island.




Figure 2. Sketch map of Connecticut, adjacent parts of New York, Rhode Island, and part of SE Massachusetts showing inferred regional SE-directed flow pattern in the glacier (No. II of our classification, Early Wisconsinan?) that deposited the Harbor Hill Moraine. (After J. H. Wilson, 1906, Plate II, fig. b, with N and E sides cropped.)

Our proposed classification of the Pleistocene deposits of the New York City region is close to Fuller's but differs in that we recognize products of five (not four) glacial advances (Table 2).

Age	Till No.	Ice-flow Direction	Description; remarks
Late Wisconsinan ("Woodfordian"?)	I	NNE to SSW	Gray-brown till in Westchester Co., Staten Is., Brooklyn, & Queens (but not present on rest of Long Island); Hamden Till in CT with terminal moraine lying along the S coast of CT; gray lake sediments at Croton Point Park, Westchester Co.
<i>Mid-Wisconsinan (?)</i>			Paleosol on Till II, SW Staten Island.
Early Wisconsinan(?)	II	NW to SE	Harbor Hill Terminal Moraine and associated outwash (Bellmore Fm. in Jones Beach subsurface); Lake Chamberlain Till in southern CT.
<i>Sangamonian(?)</i>			Wantagh Fm. (in Jones Beach subsurface).
	IIIA	NW to SE	Ronkonkoma Terminal Moraine and associated outwash (Merrick Fm. in Jones Beach subsurface).
Illinoian(?)	IIIB		Manhasset Fm. of Fuller (with middle Montauk Till Member; in lower member, coarse delta foresets (including debris flows) deposited in Proglacial Lake Long Island dammed in on S by pre-Ronkonkoma terminal moraine.
	IIIC		
<i>Yarmouthian</i>			Gardiners Clay & Jacob Sand.
Kansan(?)	IV	NNE to SSW	Gray till with decayed stones at Teller's Point (Croton Point Park, Westchester Co.); gray till with green metavolcanic stones, Target Rock, LL.
<i>Aftonian(?)</i>			No deposits; deep chemical decay of Till V.
Nebraskan (?)	V	NW to SE	Reddish-brown decayed-stone till and -outwash at AKR Co., Staten Island, and at Garvies Point, Long Island; Jameco Gravel fills subsurface valley in SW Queens.
			Pre-glacial (?) Mannelto Gravel fills subsurface valleys.

Table 2. Proposed classification of Pleistocene deposits in southeastern New York and vicinity according to our interpretation that these deposits have resulted from 5 glacial invasions, assigned roman numerals counted down from the top.

Because we lack evidence for assigning absolute ages to the products of these various glacial episodes, the best we can do is count them down from the top. Accordingly, we designate each of the tills (and its formative glacier) by a roman numeral, starting with I, the youngest, at the top, and ending with V, the oldest, at the bottom.

In this paper we present some highlights of the bedrock underlying areas north and west of Long Island that have provided distinctive indicator stones to the Pleistocene deposits and outline the components of our proposed classification.

HIGHLIGHTS OF BEDROCK UNDERLYING AREAS NORTHWEST AND NORTH OF LONG ISLAND

The bedrock underlying the regions to the northwest and north of Long Island contains vastly different assemblages including Proterozoic- to Paleozoic metamorphic rocks from the central parts of the Appalachians and Mesozoic sedimentary redbeds and intercalated mafic igneous rocks that are remnants of the fillings of the Newark and Hartford basins, respectively.

In addition, during the Cretaceous, coastal-plain strata, whose modern-day outcrop edge extends across NE New Jersey and is buried beneath the Pleistocene sediments of Long Island, extended inland for an unknown distance in southern New England. Although NW and N of the outcrop edge, most of the overlapping coastal-plain strata have been eroded, remnants are present locally at the surface and are well known in the subsurface of New York City and vicinity.

Together, the crystalline metamorphic rocks underlying the region of interest consist of sillimanite- and kyanite-grade Proterozoic- to Lower Paleozoic rocks, including gneiss, quartzite, marble, schist, and amphibolite (Baskerville, 1992, 1994; Rodgers, 1985; Drake and others, 1996). Distinctive igneous rocks of Late Ordovician age are present, notably in the Cortlandt Complex near Peekskill, Westchester Co., NY (Ratcliffe, 1981; Ratcliffe, Bender, and Tracy, 1983).

Two bodies of Mesozoic-age strata nonconformably overlie these deformed crystalline-basement rocks (Fig. 3). The Mesozoic rocks (of Late Triassic-Early Jurassic ages, Olsen, 1978; Froelich and Olsen, 1985; Lyttle and Epstein, 1987) consist predominantly red-colored sedimentary strata but include intercalated sheets of both intrusive-and extrusive dark bluish-gray mafic igneous rocks whose fresh color is masked by a thin limonitic coating from the oxidation of thin film of joint-facing pyrite and whose textures range from that of basalt to that of pegmatitic gabbro.

West of the Hudson River, from Stony Point, Rockland Co., NY, south to Staten Island and beyond, is the type area of the Newark Supergroup (Froelich and Olsen, 1985). Here, the strata dip predominantly to the NW, usually at angles of about 15° or less (Lyttle and Epstein, 1987). The tilted and eroded edge of the sheet of resistant intrusive rock near the base of the sedimentary strata forms the Palisades ridge. Comparable tilted, eroded edges of sheets of resistant volcanic rocks that are intercalated high up in the stratigraphic succession form the Watchung ridges of north-central New Jersey.

The Mesozoic sedimentary and igneous rocks filling the Hartford Basin underlie a north-south-trending lowland in the central part of Connecticut that continues northward into Massachusetts. The strata composing the filling the Hartford basin show many similarities with those filling the Newark basin in New Jersey and adjacent Pennsylvania (Sanders, 1963). Both consist predominantly of red-colored sedimentary rocks and intercalated sheets of mafic igneous rocks both intrusive and extrusive. In contrast with the Newark basin, the regional dip of the preserved remnants of the Hartford basin-filling strata is to the E. The tilted edge of the resistant intrusive sheet near the base of the Mesozoic rocks in the central Connecticut belt forms West Rock. The Connecticut equivalents of the Watchung extrusives are the volcanic rocks of the Talcott, Holyoke, and Hampden basalts. These underlie curvilinear ridges that are near the top of the preserved sedimentary strata in southern CT, but are present near the middle of the succession in central CT.




Figure 3. Regional sketch map of part of NE USA showing locations of preserved remnants of strata that filled Mesozoic nonmarine basins (Newark basin in NY, NJ, and PA; Hartford basin in CT and MA; light gray tone). Parallel line segments show how glacial ice flowing regionally from each of the observed dominant directions would distribute diagnostic erratics to Long Island. P = Peekskill, NY, near unique Cortlandt Complex of coarse ultramafic igneous rocks. (Base map from J. E. Sanders, 1963, fig. 7, p. 513.)

A. Flow from NNW to SSE deposits reddish-brown till and erratics of Mesozoic rocks in extreme W and extreme E Long Island, but throughout a substantial stretch along the N shore of Long Island, downflow from the "crystalline corridor" between the two belts of Mesozoic rocks, did not deposit any reddish-brown till. The distribution of erratics in the Harbor Hill Moraine and in the Ronkonkoma Moraine proves these moraines were deposited by ice flow from NNW to SSE, as shown in Fig. 2.

B. Flow from NNE to SSW, as in the "Woodfordian" glacier, distributes erratics from the Hartford basin to the SSW. The absence of reddish-brown till and associated Mesozoic erratics in the Pleistocene deposits exposed along the N shore of Long Island WSW of the S end of the Hartford basin proves that the "Woodfordian" ice could not have deposited the main bodies of Long Island's two famous terminal-moraine ridges.

The distinctive colors and compositions of the Mesozoic rocks make them ideal indicator stones and pigments for distinctive reddish-brown tills. However, by themselves, erratics from the Newark basin cannot be distinguished from those from the Hartford basin. At least locally, however, the associated non-Mesozoic rocks provide useful diagnostic clues. For example, low-grade phyllites, -schist, and -metavolcanic rocks containing chlorite and epidote are present in the western Connecticut terrane W of the S end of the Hartford basin (Fritts, 1962; Burger, 1967). No such low-grade rocks of volcanoclastic parentage are present in the vicinity of the Newark Basin of New Jersey. Therefore, erratics of such low-grade volcanoclastic rocks associated with distinctive Mesozoic rocks would **uniquely** identify a Connecticut source.

COMPONENTS OF OUR PROPOSED CLASSIFICATION

OF PLEISTOCENE DEPOSITS

As mentioned, from our previous work (summarized in Sanders and Merguerian, 1997) we know that in New York City, glacial sediments deposited by ice flowing from NNW to SSE (**across** the Hudson Valley) are characterized by their distinctive reddish-brown-color, the result of a glacier's grinding over hematite-rich sedimentary rocks from the Newark Basin. By contrast, sediments deposited by glaciers that flowed from the NNE to the SSW (**down** the Hudson Valley) are associated with yellow-brown- to brownish-gray tills, the result of glacial scour of non-hematite-bearing rocks underlying the "crystalline corridor" of metamorphic rocks exposed between the Newark- and Hartford basins. Each of the flow directions resulted in a system of crosscutting glacial features and diagnostic indicator stones, easily identifiable in the field.

According to us, **Glacier I** flowed from NNE to SSW and did not reach most of Long Island; it deposited the fields of drumlins in Rockland County, NY and the yellowish-brown till forming a single drumlin at Enoch's Nose, Croton Point Park, Westchester County, NY; the Hamden Till in south-central Connecticut (Flint, 1961); and a terminal moraine along the S coast of Connecticut [SW of the Norwalk Islands (Ellis, 1962), it crossed what is now Long Island Sound and covered Queens (Woodworth, 1901; Brooklyn (Lyell, 1852), and at least some of Staten Island]. Based on evidence from the Catskills (Rich, 1935), we can infer that the "Woodfordian" glacier was thinner than most, if not all, of its predecessors (top at alt. 3900 ft vs. ice thick enough to cover all of the Catskills and all other high parts of New England). A possible Glacier I/II interglacial deposit is the paleosol capping the reddish-brown till in the coastal bluffs of SW Staten Island.

Glacier II, flowing from NNW to SSE, reached all of Long Island and deposited the Harbor Hill Moraine and reddish-brown till at Croton Point Park and other localities on the E side of the Hudson River in Westchester County and New York City; and possibly also the Lake Chamberlain Till of south-central Connecticut (Flint, 1961). Glacier II/III interglacial marginal-marine sediments include the Wantagh Formation of the Jones Beach subsurface [Rampino (1978 ms.); Rampino and Sanders (1981)].

Glacier III, the most-extensive of them all, also flowed from NNW to SSE and featured three fluctuations; its earliest advance deposited a now-vanished and -submerged terminal moraine; the ice front then retreated and a regionally extensive proglacial Lake Long Island formed in which the lacustrine sediments forming the lower member of Fuller's Manhasset Formation were deposited; a subsequent readvance deposited the Montauk Till; and after another retreat and deposition of outwash sediments, a final readvance deposited the Ronkonkoma Moraine. Glacier III/IV interglacial deposits include the Gardiners Clay.

Glacier II and/or Glacier III (we do not know which; it could have been both) brought anthracite erratics from NE Pennsylvania into the New York City region. Such ice flow explains the results that Zen and Mamay (1968), who accepted the consensus view of only a single direction of Pleistocene ice flow from the NNE to the SSW, found so puzzling.

Glacier IV flowed from the NNE to the SSW, deposited a gray-brown till exposed at water's edge, Teller's Point, Croton Point Park and in the lower part of the coastal bluff at Target Rock National Wildlife Refuge, Long Island; and sculpted many rock exposures in the Hudson Highlands, in northern Manhattan, and in the New York Botanical Garden and sculpted the roche moutonnée structure at the N end of Pelham Bay Park, both in The Bronx. We have not found any Glacier IV/V interglacial deposits.

Glacier V flowed from NNW to SSE and deposited the much-decomposed reddish-brown tills resting on the Upper Cretaceous coastal-plain strata at the AKR Excavating Company, SW Staten Island, and at Garvies Point, Long Island.

CONCLUSIONS

The main bodies of Long Island's two famous terminal moraines contain unmistakable provenance evidence that they were deposited by ice that flowed regionally from NNW to SSE. Because the "Woodfordian" glacier, our No. I, is the glacier most geologists accept as being responsible for these two moraines, flowed from NNE to SSW and stopped along the S coast of CT; it reached only westernmost Long Island (Queens and Brooklyn) and adjacent parts of Staten Island. Accordingly, the main body of the Harbor Hill Moraine **could not have been deposited by the "Woodfordian" glacier** but rather must have been deposited by ice flowing from NNW to SSE. We think the glacier responsible for depositing the main body of the Harbor Hill Moraine was our No. II (of probable Early Wisconsinan age, as Fuller believed). The subsurface Wantagh Formation, a marginal-marine deposit, may be of Sangamonian age. The most-extensive local Pleistocene unit is Fuller's Manhasset Formation, which is probably of Illinoian age. It consists of two deposits of large proglacial lakes separated by the Montauk Till (our units IIIA, IIIB, and IIIC). We conclude that a terminal moraine of pre-Ronkonkoma age (deposited by our Glacier IIIC), which we correlate with the Martha's Vineyard-Nantucket moraines and bouldery shoals in between, served as the dam to impound Proglacial Lake Long Island in which the lacustrine parts of the Manhasset Formation were deposited. We accept Wehmiller's finding that the Gardiners Clay is of Yarmouthian age (as assigned by Fuller) and **not of Sangamonian age** as supposed by MacClintock and Richards (1936) and their host of followers. The ages of two pre-"Illinoian" tills (our Nos. IV and V) that contain much-decayed stones are not well constrained. We guesstimate their assignments as "Kansan" (?) and "Nebraskan" (?), respectively.

ACKNOWLEDGEMENTS

Our continued studies of the glacial geology of the New York City area have been supported by the Geology Department of Hofstra University. We are indebted to our colleague Prof. J Bret Bennington for his interest and observations made on several joint field trips.

REFERENCES CITED

- Baskerville, C. A., 1992, Bedrock (*sic*) and engineering geologic (*sic*) maps of Bronx County and parts of New York and Queens counties, New York: U. S. Geological Survey Miscellaneous Investigations Series Map I-2003 (scale 1:24,000).
- Baskerville, C. A., 1994, Bedrock (*sic*) and engineering geology maps of New York County and parts of Kings and Queens counties, New York and parts of Bergen and Hudson counties, New Jersey: U. S. Geological Survey Miscellaneous Investigations Series Map I-2306 (2 sheets; colored maps on scale of 1/24,000).
- Burger, H. R., III, 1967, Stratigraphy and structure of the western part of the New Haven quadrangle, Connecticut: Connecticut Geological and Natural History Survey Report of Investigations 4, 15 p. (geologic map on scale of 1/57,600).
- Cadwell, D. H., *ed.*, 1986, The Wisconsinan Stage of the First Geological District, eastern New York: New York State Museum, Bulletin 455, 191 p.
- Cadwell, D. H., 1989, Surficial geology map of New York, Lower Hudson sheet: New York State Museum Map and Chart Series 40, scale 1:250,000.
- Chamberlin, T. C., 1883, Preliminary paper on the terminal moraine of the second glacial epoch, p. 291-402 *in* United States Geological Survey Annual Report, 3rd (1881-1882), 564 p. (1843-1928)
- Chamberlin, T. C., 1888, The rock scorings of the great Ice Age: U. S. Geological Survey Annual Report, 7th (1885-1886), p. 147-248.
- Chamberlin, T. C., 1895a, The classification of American glacial deposits: *Journal of Geology*, v. 3, p. 270-277.
- Chamberlin, T. C., 1895b, Glacial phenomena of North America, p. 724-775 *in* Geikie, James, The great ice age: New York, D. Appleton and Company, 000 p.
- Clapp, F. G., 1908, Complexity of the glacial period in northeastern New England: *Geological Society of America Bulletin*, v. 18, p. 505-556.
- Crosby, W. O., 1890, Geological history of the Boston basin [Mass.]: *Boston Society of Natural History Proceedings*, v. 25, p. 10-17.
- Crosby, W. O., 1891, Composition of the till or boulder clay: *Boston Society of Natural History Proceedings*, v. 25, p. 115-140.
- Drake, A. A., Volkert, R. A., Monteverde, D. H., Herman, G. C., Houghton, H. F., Parker, R. A., and Dalton, R. F., 1996, Bedrock geologic map of northern New Jersey: U. S. Geological Survey Miscellaneous Map Investigations I-2540-A (colored geologic map in 2 sheets, scale 1/100,000).
- Ellis, C. W., 1962, Marine sedimentary environments in the vicinity of the Norwalk Islands, Connecticut: *Connecticut Geological and Natural History Survey Bulletin* 94, 89 p.
- Fleming, W. L. S., 1935, Glacial geology of central Long Island: *American Journal of Science*, 5th series, v. 30, p. 216-238.
- Flint, R. F., 1935, How many glacial stages are recorded in New England?: *Journal of Geology*, v. 43, p. 771-777.
- Flint, R. F., 1945, Glacial map of North America: *Geological Society of America Special Papers* No. 60, 37 p.
- Flint, R. F., 1961, Two tills in southern Connecticut: *Geological Society of America Bulletin*, v. 72, no. 11, p. 1687-1691.
- Flint, R. F., and Gebert, J. A., 1976, Latest Laurentide ice sheet: new evidence from southern New England: *Geological Society of America Bulletin*, v. 87, no. 2, p. 182-188.
- Fritts, C. E., 1962, Age (*sic*) and sequence of metasedimentary (*sic*) and metavolcanic formations northwest of New Haven Connecticut: U. S. Geological Survey Professional Paper 450-D, p. D32-D36.
- Froelich, A. J., and Olsen, P. E., 1985, 1. Newark Supergroup, a revision of the Newark Group in eastern North America, p. 1-3 *in* Robinson, G. R., and Froelich, A. J., *eds.*, U. S. Geological Survey workshop on the Early Mesozoic basins of the eastern United States, Second, Proceedings: U. S. Geological Survey Circular 946, 147 p.
- Fuller, M. L., 1901, Probable representatives of pre-Wisconsin till in southeastern Massachusetts: *Journal of Geology*, v. 9, p. 311-329.
- Fuller, M. L., 1903, Probable pre-Kansan and Iowan deposits of Long Island: *American Geologist*, v. 32, p. 308-312.
- Fuller, M. L., 1905, Geology of Fishers Island, New York: *Geological Society of America Bulletin*, v. 16, p. 367-390. [Gardiner (*sic*; no s to make Gardiners) clay on 375-378]

- Fuller, M. L., 1906, Glacial stages in southeastern New England and vicinity: *Science*, new series, v. 24, p. 467-469. considers that the till at Brockton is pre-Kansan; that the Boston drumlins are composed of Illinoian till, and the thin surficial till is the Wisconsin (total: 3).
- Fuller, M. L., 1914, The geology of Long Island, New York: U. S. Geological Survey Professional Paper 82, 223 p.
- Fuller, M. L., 1937, Comment on: "Correlation of Late (*sic*) Pleistocene marine (*sic*) and glacial deposits of New Jersey and New York:" *Geological Society of America Bulletin*, v. 47 Supplement, p. 1982-1994.
- Gilbert, G. K., 1885, The topographic features of lake shores, p. 69-123 in U. S. Geological Survey, Annual Report, 5th (1883-1884), 469 p.
- Gilbert, G. K., 1890, Lake Bonneville: U. S. Geological Survey, Monograph 1, 438 p.
- Goldsmith, Richard, 1960, A post Harbor Hill-Charlestown moraine in southeastern Connecticut: *American Journal of Science*, v. 258, p. 740-743.
- Isachsen, Y. W.; Landing, Ed; Lauber, J. M.; Rickard, L. V.; and Rogers, W. B., *eds.*, 1991, *Geology of New York. A simplified account*: New York State Museum and Science Service Educational Leaflet 20, 284 p.
- Kaye, C. A., 1964a, Boulder train of silicified Paleozoic wood, southeastern Massachusetts: *Geological Society of America Bulletin*, v. 75, no. 3, p. 233-236.
- Kaye, C. A., 1964b, Outline of Pleistocene geology of Martha's Vineyard, Massachusetts: United States Geological Survey Professional Paper 501-C, p. C134-C139.
- Kaye, C. A., 1964c, Illinoian and early Wisconsin moraines of Martha's Vineyard, Massachusetts: United States Geological Survey Professional Paper 501-C, p. C140-C143.
- Kaye, C. A., 1964d, The Pleistocene geology of Martha's Vineyard, Massachusetts: Itinerary for the Friends of the Pleistocene Annual Reunion, 27th: U. S. Geological Survey Open-File Report.
- Kaye, C. A., 1976, The geology (*sic*) and early history of the Boston, Massachusetts area--a bicentennial approach: *United States Geological Survey Bulletin* 1476, 78 p.
- Kaye, C. A., 1982, Bedrock (*sic*) and Quaternary geology of the Boston area, Massachusetts, p. 25-40 in Legget, R. F., *ed.*, *Geology under cities: Boulder, Colorado*, The Geological Society of America, *Reviews in Engineering Geology*, v. 5, 131 p.
- Koteff, Carl; Oldale, R. N.; and Hartshorn, J. H., 1967, Geologic map of the North Truro quadrangle, Barnstable County, Massachusetts: U. S. Geological Survey Geologic Quadrangle Map GQ-599, scale 1/24,000.
- Lyell, Sir Charles, 1852, *Travels in North America, in the years 1841-2; with geological observations on the United States, Canada, and Nova Scotia*: New York, NY, John Wiley, 2 volumes. (v. 1, 251 p.; v. 2, 197 p.) (according to Muller, 1965, p. 48, the date is 1845 and 221 p in v. 2)
- Lytle, P. T., and Epstein, J. B., 1987, Geologic map of the Newark 1° by 2° quadrangle, New Jersey, Pennsylvania, and New York: United States Geological Survey Miscellaneous Investigations Series Maps, Map I-1715 (2 sheets; scale 1:250,000).
- MacClintock, Paul; and Richards, H. G., 1936, Correlation of the late (*sic*) Pleistocene marine (*sic*) and glacial deposits of New Jersey and New York: *Geological Society of America Bulletin*, v. 47, no. 3, p. 289-338.
- MacClintock, Paul; and Richards, H. G., 1937, Correlation of the late (*sic*) Pleistocene marine (*sic*) and glacial deposits of New Jersey and New York (reply to Discussion by M. L. Fuller): *Geological Society of America Bulletin*, v. 47 Supplement, p. 1982-1994.
- Muller, E. H., 1964, Quaternary geology of New York, p. 99-112 in Wright, H. E., and Frey, D. G., *eds.*, *The Quaternary of the United States*: Princeton, NJ, Princeton University Press, 922 p.
- Muller, E. H., and Calkin, P. E., 1993, Timing of Pleistocene glacial events in New York State: *Canadian Journal of Earth Sciences*, v. 30, p. 1829-1845.
- Muller, E. H.; Sirkin, Les; and Craft, J. L., 1993, Stratigraphic evidence of a pre-Wisconsinan interglaciation in the Adirondack Mountains, New York: *Quaternary Research*, v. 40, p. 163-138.
- Newman, W. A., and Mickelson, D. M., 1994, Genesis of Boston Harbor drumlins, Massachusetts, p. 333-343 in Dardis, G. F., and McCabe, A. M., *eds.*, *Subglacial processes, sediments (*sic*) and landforms (*sic*): Symposium, University of Ulster, July 1992: *Sedimentary Geology*, v. 91, nos. 1-4 (special volume), 382 p.*
- Oldale, R. N., 1982, Pleistocene stratigraphy of Nantucket, Martha's Vineyard, The Elizabeth Islands, and Cape Cod, Massachusetts, p. 1-34 in Larson, G. J., and Stone, B. D., *eds.*, *Late Wisconsinan glaciation of New England. Proceedings of a symposium held at Northeastern section of Geological Society of America, 13 March 1980, Philadelphia, PA. Dubuque, Iowa, Kendall/Hunt Publishing Company, 242 p.*

- Oldale, R. N., and Colman, S. M., 1992, On the age of the penultimate full glaciation of New England, p. 163-170 in Clark, P. U., and Lea, P. D., eds., *The last Interglacial-Glacial transition in North America*: Boulder, CO, Geological Society of America Special Paper 270, 317 p.
- Oldale, R. N., and Eskenasy, D. M., 1983, Regional significance of pre-Wisconsin till from Nantucket Island, Massachusetts: *Quaternary Research*, v. 19, p. 302-311.
- Oldale, R. N., Valentine, P. C., Cronin, T. M., Spiker, E. C., Blackwelder, B. W., Belknap, D. F., Wehmiller, J. F., and Szabo, B. J., 1982, Stratigraphy (*sic*), structure (*sic*), absolute age (*sic*), and paleontology of the upper Pleistocene deposits at Sankaty Head, Nantucket Island, Massachusetts: *Geology*, v. 10, no. 5, p. 246-252.
- Olsen, P. E., 1978, On the use of the term Newark for Triassic and Early (*sic*) Jurassic rocks of eastern North America: *Newsletters Stratigraphy*, v. 7, no. 2, p. 90-95.
- Pessl, Fred, Jr., 1966, A two-till locality in northeastern Connecticut: U. S. Geological Survey Professional Paper 550-D, P. D89-D93.
- Pessl, Fred, Jr.; and Schafer, J. P., 1968, Two-till problem in Naugatuck-Torrington area, western Connecticut, Trip B-1. p. B1-B25 in Orville, P. M., ed., *Guidebook for field trips in Connecticut*: New England Intercollegiate Geological Conference Annual Meeting, 62nd, New Haven, Connecticut: Hartford, CT, Connecticut Geological and Natural History Survey Guidebook No. 2, not consecutively paginated.
- Rampino, M. R., 1978 ms., Quaternary history of south-central Long Island, New York: New York, New York, Columbia University, Department of Geological Sciences, Ph. D. Dissertation, 742 p.
- Rampino, M. R., and Sanders, J. E., 1981, Upper Quaternary stratigraphy of southern Long Island, New York: *Northeastern Geology*, v. 3, no. 2, p. 116-128.
- Ratcliffe, N. M., 1981, Cortlandt-Beemerville magmatic belt: a probable late Taconian alkalic cross trend in the central Appalachians: *Geology*, v. 9, no. 7, p. 329-335.
- Ratcliffe, N. M., Bender, J. F., and Tracy, R. J., 1983, Tectonic setting (*sic*), chemical petrology (*sic*) and petrogenesis of the Cortlandt Complex and related igneous rocks of southeastern New York State: Geological Society of America Northeastern Section Meeting, 23-26 March 1983, Concord Hotel, Kiamesha Lake, NY, Guidebook for Field Trip 1, 101 p.
- Rich, J. L., 1935, Glacial geology of the Catskills: *New York State Museum Bulletin No. 299*, 180 p.
- Ricketts, H. C., 1986, Examination in boreholes of the subsurface Gardiners Clay (Pleistocene), in the context of the Cretaceous-Quaternary section of Great Neck, Long Island, New York: *Northeastern Geology*, v. 8, nos. 1/2, p. 13-22.
- Rodgers, John, 1985, *Bedrock geological map of Connecticut*: Hartford, CT, Connecticut Geological and Natural History Survey, Connecticut Natural Resources Atlas Series, scale 1:250,000.
- Salisbury, R. D., assisted by H. B. Kümmel, C. E. Peet, and G. N. Knapp, 1902, *The glacial geology of New Jersey*: Geological Survey of New Jersey, Final Report of the State Geologist, v. 5: Trenton, New Jersey, MacCrellish and Quigley, Book and Job Printers, 802 p.
- Salisbury, R. D., and Peet, C. E., 1894, Section V. Drift phenomena of the Palisade ridge, p. 157-224 in *New Jersey State Geologist Annual Report for 1893*: Trenton, NJ, The John L. Murphy Publishing Company Printers, 457 p.
- Sanders, J. E., 1963, Late Triassic tectonic history of northeastern United States: *American Journal of Science*, v. 261, p. 501-524.
- Sanders, J. E., 1974, Geomorphology of the Hudson Estuary, p. 5-38 in Roels, Oswald, ed., *Hudson River Colloquium*: New York Academy of Sciences Annals, v. 250, 185 p.
- Sanders, J. E.; and Merguerian, Charles, 1991, Pleistocene geology of Long Island's north shore: Sands Point and Garvies Point to Target Rock: Long Island Geologists' Association Field Trip 29 June 1991 Guidebook: Hempstead, NY, Hofstra University Department of Geology, 40 p.
- Sanders, J. E.; and Merguerian, Charles, 1994a, Fitting newly discovered north-shore Gilbert-type lacustrine deltas into a revised Pleistocene chronology of Long Island (extended abstract), p. 103-113 in Hanson, G. N., *chm.*, *Geology of Long Island and metropolitan New York*, 23 April 1994, Stony Brook, NY: Stony Brook, NY, Long Island Geologists Program with Abstracts, 165 p.
- Sanders, J. E.; and Merguerian, Charles, 1994b, Glacial geology of the New York City region, p. 93-200 in Benimoff, A. I., ed., *The geology of Staten Island*, New York: Geological Association of New Jersey Annual Meeting, 11th, Somerset, NJ, 14-15 October 1994, Field guide and proceedings, 296 p.
- Sanders, J. E.; and Merguerian, Charles, 1995, Evidence for pre-Woodfordian ages of Long Island's terminal moraines, p. 91-106 in Hanson, G. N., *chm.*, *Geology of Long Island and metropolitan New York*, 22 April 1995, Stony Brook, NY: Stony Brook, NY, Long Island Geologists Program with Abstracts, 135 p.
- Sanders, J. E.; and Merguerian, Charles, 1996, Trip 39: Glacial geology of Long Island, 01-02 June 1996 (revision of Trip 15, 17-18 November 1990): New York Academy of Sciences Section of Geological Sciences Trips on the Rocks Guidebook, 129 p.

- Sanders, J. E.; and Merguerian, Charles, 1997, Geologic setting of a cruise from the mouth of the East River to the George Washington bridge, New York Harbor: New York, NY, American Rock Mechanics Association, NYRocks 97, hosted by Columbia University Field-Trip Guidebook, 147 p.
- Sanders, J. E.; Merguerian, Charles; and Mills, H. C., 1993, "Port Washington deltas" of Woodworth (1901) revisited: pre-Woodfordian Gilbert-type deltas revealed in storm-eroded coastal bluff, Sands Point, New York (abstract): Geological Society of America Abstracts with Programs, v. 25, no. 6, p. A-308.
- Sanders, J. E.; Merguerian, Charles; and Okulewicz, S. C., 1995, Recumbent fold in displaced slab of Upper Cretaceous sediments, Princes Bay, Staten Island, New York: Further evidence that ice flowing southeastward deposited the Harbor Hill Moraine (extended abstract), p. 107-117 in Hanson, G. N., *chm.*, Geology of Long Island and metropolitan New York, 22 April 1995, Stony Brook, NY: Stony Brook, NY, Long Island Geologists Program with Abstracts, 135 p.
- Seyfert, C. K., and Leveson, D. J., 1968, Structure (*sic*) and petrology of Pelham Bay Park, p. 175-195 in Finks, R. M., *ed.*, New York State Geological Association Annual Meeting, 40th, Queens College, Flushing, New York, Guidebook to Field Excursions: Flushing, NY, Queens College of the City University of New York Department of Geology, 253 p.
- Schafer, J. P., and Hartshorn, J. H., 1965, The Quaternary of New England, p. 113-128 in Wright, H. E., Jr.; and Frey, D. C., *eds.*, The Quaternary of the United States: Princeton, NJ, Princeton University Press, 922 p.
- Schuberth, C. J., 1968, The geology of New York City and environs: Garden City, NY, Natural History Press, 304 p.
- Shaler, N. S., 1866, Notes on the position and character (*sic*) of some glacial beds containing fossils at Gloucester, Massachusetts: Boston Society of Natural History Proceedings, v. 11, p. 27-30. (till overlying fossiliferous sediments at Gloucester, MA.)
- Shaler, N. S., 1889b, The geology of Nantucket: U. S. Geological Survey Bulletin 53, 55 p.
- Shaler, N. S., Woodworth, J. B., and Marbut, C. F., 1896, The glacial brick clays of Rhode Island and southeastern Massachusetts, p. 951-1004 in U. S. Geological Survey Annual Report, 17th, part 1.
- Sirkin, L. A., 1968, Geology (*sic*), geomorphology (*sic*) and late glacial (*sic*) environments of western Long Island, New York, p. 233-253 in Finks, R. M., *ed.*, New York State Geological Association Annual Meeting, 40th, Queens College, Guidebook to field trips: Flushing, NY, Queens College of City University of New York Department of Geology, 253 p.
- Sirkin, L. A., 1982, Wisconsinan glaciation of Long Island, New York to Block Island, Rhode Island, p. 35-59 in Larson, G. J., and Stone, B. D., *eds.*, Late Wisconsinan glaciation of New England. Proceedings of a symposium held at Northeastern section of Geological Society of America, 13 March 1980, Philadelphia, PA. Dubuque, Iowa, Kendall/Hunt Publishing Company, 242 p.
- Sirkin, L. A., 1986, Pleistocene stratigraphy of Long Island, New York, p. 6-21 in Cadwell, D. H., *ed.*, The Wisconsinan Stage of the First Geological District, eastern New York: New York State Museum Bulletin 455, 191 p.
- Sirkin, L. A., 1991, Stratigraphy of the Long Island platform, p. 217-227 in Gayes, P., Lewis, R. S., and Bokuniewicz, H. J., *eds.*, Quaternary geology of Long Island Sound: Journal of Coastal Research, Walter Newman Memorial volume, Special Issue No. 11, 215 p.
- Sirkin, L. A., 1994, Geology of the Long Island platform (extended abstract), p. 137-139 in Hanson, G. N., *chm.*, Geology of Long Island and metropolitan New York, 23 April 1994, Stony Brook, NY: Stony Brook, NY, Long Island Geologists Program with Abstracts, 165 p.
- Sirkin, L. A., 1995a, Deglaciation of eastern Long Island: The terminal moraine, recessional moraines, outwash plains, proglacial lakes and meltwater channels (extended abstract), p. 127-134 in Hanson, G. N., *chm.*, Geology of Long Island and metropolitan New York, 22 April 1995: Stony Brook, NY: Stony Brook, NY, Long Island Geologists Program with Abstracts, 135 p.
- Sirkin, L. A., 1995b, Eastern Long Island geology with field trips: Watch Hill, RI, Book and Tackle Shop, 220 p.
- Sirkin, L. A., and Mills, H. C., 1975, Wisconsinan glacial stratigraphy and structure of northwestern Long Island, Trip B5, p. 299-327 in Wolff, M. P., *ed.*, Guidebook to Field Excursions: New York State Geological Association, Annual Meeting, 47th, Hofstra University, Hempstead, New York: Hempstead, N.Y., Hofstra University, Department of Geology, 327 p.
- Sirkin, L. A.; and Stuckenrath, R., 1980, The Portwashingtonian warm interval in the northern Atlantic coastal plain: Geological Society of America Bulletin, v. 91, p. 332-336.
- Stevens, R. P., 1872, On glacial phenomena in the vicinity of New York City: American Journal of Science, 3rd series, v. 4, p. 88-90.
- Upham, Warren, 1879a, Terminal moraines of the North American ice sheet: American Journal of Science, 3rd series, v. 18, p. 81-92.
- Upham, Warren, 1879b, Glacial drift in Boston and its vicinity [Mass.]: Boston Society of Natural History Proceedings, v. 20, p. 220-234.
- Upham, Warren, 1880, The succession of glacial deposits in New England: American Association Proceedings, v. 28, p. 299-310.

Upham, Warren, 1889a, Glaciation of mountains in New England and New York: Appalachia, v. 5, p. 291-312. (Also, American Geologist, v. 4, p. 165-174.)

Upham, Warren, 1889b, Marine shells and the fragments of shells in the till near Boston: Boston Society of Natural History Proceedings, v. 24, p. 127-141.

Upham, Warren, 1889c, The structure of drumlins: Boston Society of Natural History Proceedings, v. 24, p. 228-242.

Widmer, Kemble, 1964, The geology (*sic*) and geography of New Jersey: The New Jersey Historical Series, v. 19: Princeton-New York-Toronto-London, D. Van Nostrand Company, Inc., 193 p.

Wilson, J. H., 1906, The glacial history of Nantucket and Cape Cod with an argument for a fourth center of glacial dispersion in North America: New York, NY, Columbia University Press, 90 p.

Woodworth, J. B., 1901, Pleistocene geology of portions of Nassau County and Borough of Queens (N.Y.): New York State Museum Bulletin 48, p. 618-670. (Includes colored geologic map on scale of 1:62,500.)

Woodworth, J. B.; and Wigglesworth, Edward, *eds.*, 1934, Geography (*sic*) and geology of the region including Cape Cod, the Elizabeth Islands, Nantucket, Martha's Vineyard, No Man's Land, and Block Island: Cambridge, MA, Harvard College Museum of Comparative Zoology Memoir, v. 52, 338 p.

Zen, E-An; and Mamay, S. H., 1968, Middle Pennsylvanian plant fossils: problematic occurrence in the Bronx: Science, v. 161, p. 157-158.

This Reference:

Sanders, John E., and Merguerian, Charles, 1998, Classification of Pleistocene deposits, New York City and vicinity – Fuller (1914) revived and revised: p. 130-143 in Hanson, G. N., *chm.*, Geology of Long Island and Metropolitan New York, 18 April 1998, State University of New York at Stony Brook, NY, Long Island Geologists Program with Abstracts, 161 p.

Filename: JESCM98.pdf