American Society of Civil Engineers


Evaluating Geological Controls on Hard Rock Excavation, New York City, NY


Charles Merguerian

















~ 450 Ma Taconic Arc – Passive Margin Collision







450 Ma to 250 Ma Protracted Plate Collisions Form the Appalachians








Merguerian Has Spent Most of his Career Mapping the Surface and Subsurface Geology of NYC

> Proper Field Attire For NYC

O-Ch Hartland Formation and Manhattan Schist (upper unit) WESTCHESTER COUNTY Omm Manhattan Schist (middle unit) Oml Manhattan Schist 0-Eh (lower unit) 0-Ei Inwood Marble HARTLAND TERRANE Fordham Gneiss (HUTCHINSON RIVER GROUP Yonkers Gneiss 0-€h Long Island Sound River 0-€ East HARTLAND TERRANE 0-Eh

Hudson River O-Ch St. Nicholas thrust 0 0.5 1 2 KILOMETER

New York City





Merguerian, 2001













Nalloomsac "Balmville" Contact, Grand Concourse, Bronx, Ny







Manhattan Schist F₃ Folds of S₂ Central Park, NYC















O-Ch Hartland Formation and Manhattan Schist (upper unit) WESTCHESTER COUNTY Omm Manhattan Schist (middle unit) Oml Manhattan Schist 0-Eh (lower unit) 0-Ei Inwood Marble HARTLAND TERRANE Fordham Gneiss (HUTCHINSON RIVER GROUP Yonkers Gneiss 0-€h Long Island Sound River 0-€ East HARTLAND TERRANE 0-Eh


Hudson River O-Ch St. Nicholas thrust 0 0.5 1 2 KILOMETER

New York City



Merguerian, 2001







- Published Maps and Reports Boring Analysis Fractures **Rock Types Rock Fabrics Density Studies Petrographic Studies** Rock Fabric Studies
 - Mineralogy and Texture
 Structure
 Orientation

Metamorphism

What Are the Geological Controls on Effective Hard Rock TBM Tunneling in Crystalline Terrains?

Excessive Fines
Blocky Ground
Unstable Headings and Sidewalls
Low Penetration Rates










Unforseen Tunneling Problems





Anticipated vs. Actual Penetration Rate










Merguerian's Queens Tunnel Field Office

Comparative Lithologic Analysis




Density Queens Tunnel (Mean = 2.87 g/cm³)







Unexpected High Garnet Content


- •The boring logs cite the term garnetiferous throughout. To most geologists, "garnetiferous" rocks contain a few % garnet
- •Thirty two Queens Tunnel Garnet Zones mapped. They underlie 2,663' or 10.64% of as-built tunnel
- •The Queens Tunnel rocks contain up to 50% garnet
- •The Queens Tunnel Garnet Concentrations would be called "ore deposits" in many parts of the world
- •Results in abrasivity to cutters and production of excessive fines








Granulite Facies Gneisses
Found in the Queens Tunnel
and Elsewhere With
Granoblastic Textures =
Tough Rocks for Excavation!











SE Manhattan






Five Localities
South of
Canal Street


Display
Sheared
Manhattan,
Hartland, and
Walloomsac
Rocks!


after Baskerville 1994, Merguerian and Moss 2007



of Berkey (1910)


Merguerian, 1984







Interpretive NW-SE Geologic Section Based on Berkey 1910 Boring Data

Merguerian, 1984






Interpretive NW-SE Geologic Section Based on Berkey 1910 Boring Data




In Western and Central Manhattan:
Amphibolite Facies Schists Well-layered Hartland Fm.
Penetrative Foliated Textures
Great Rocks for Tunneling and Excavation!












Low Mica Content 16%

Granofels

Interlocking Texture is "Knitted" by Micas

Stretched Garnet and Discontinuous Folia = Polymetamorphism




Low Mica Content 22%

Folded Migmatite Gneiss

Interlocking Texture is "Knitted" by Micas Grown at Many Angles

Discontinuous Foliation = Polymetamorphism N753




Low Mica Content 31%

Migmatite Gneiss

Coarse Micas


Discontinuous Folia




Mica Content of Rock Fabric


Micaceous (+/- hornblende) penetrative foliation

vs. non-foliated "granoblastic" rock mass






Foliated = Schist




Non-Foliated = Gneiss

Structural Fabrics




















Low Mica Content <50%

Structurally Folded Multi-directional Fabric

Interlocking Texture is "Knitted" by Micas

Polymetamorphism





N701E

Southern Manhattan Hartland Gneiss and Granofels Unit

- Low Mica Content (26%) +
 - **Granoblastic Textures +**
 - **Interlocking Minerals +**
 - Discontinuous Fabric +
 - **Variable Orientation =**

Poor Excavation Rates





HERRENKNECHT TBM ADVERTISEMENT

Geology

Rock Does Not

Equal Rock

At Herrenknecht, maximum tunnelling performance and the greatest possible safety are the ultimate goals for the development of tunnelling machines. Expert analysis of the geological conditions result in a "tailored hard rock machine".


Whether it is solid rock, abrasive rock or rock under high pressure, weathered transition zones with high ground water pressure or caverns, the variety of the geological conditions in a planned tunnel route is virtually unlimited. At Herrenknecht, the geological analysis

of the ground conditions is always taken into consideration in the machine design. Cutters and cutterhead are ideally adapted to the varying degrees of hardness and abrasion in sedimentary, metamorphic or igneous rock.

All Experts On One Team

The excavation process in hard rock takes place in the peak state of the shear and compression resistance as well as tensile strengths of the rock. At the same time, the best possible tunnelling performance has to be achieved. ping the machine design.

At Herrenknecht, a team of internal specialists from the disciplines of rock mechanics, mechanical engineering and process technology find the optimum project solution for develo-



Mechanical rock excavation is confronted by rock with varying degrees of hardness, e.g. with extremely hard gneiss (top left) and granite (top right) medium hard mica schist (center left), breccia (center right) and claystone (bottom left) as well as limestone (bottom right).





The formation of each mountain range is unique. Lötschberg in Switzerland consists of a wide variety of rock formations along the tunnel route. Herrenknecht supplied two single gripper machines (Ø 9.43 m), which enable mechanical rock stabilization as close as 4.2 m behind the cutterhead.



